

Characterization of protein-ligand complexes using molecular dynamics simulations by consideration of GABA_A protein as receptor and Lavender compounds as ligand

¹Ali Kazemi Babaheydari,^{2,*} Somayeh Faraji

¹Department of Physical Chemistry, Islamic Azad University Shahrekord Branch, Shahrekord, Iran

² Department of Physics, Islamic Azad University Shahrekord Branch, Shahrekord, Iran

*Corresponding Author: sfaraji_ph@yahoo.com; kazemibali@gmail.com

Abstract:

GABA_A receptors are a class of receptors that respond to the neurotransmitter gamma-aminobutyric acid (GABA), the chief inhibitory neurotransmitter in the vertebrate central nervous system, which are ligand-gated ion channels and are known as ionotropic receptors [1-5]. Lavender is a species of flowering plants in the mint family and is native to the mountainous zones of the Mediterranean and flourishes throughout southern Europe, Australia, and the United States. This herb has been used as a remedy for a range of ailments from insomnia and anxiety to depression and fatigue. This plant is grown mainly for the production of lavender's oil which has antiseptic and anti-inflammatory properties. A number of studies have reported that this oil may be beneficial in a variety of conditions, including insomnia, hair loss, anxiety, stress, and postoperative pain [6, 7]. This plant is being studied for antibacterial and antiviral properties [8]. In this work, we have simulated and studied the complexes of different chemical compounds found in the lavender extract with GABA protein as receptor. Our Simulations are based on molecular dynamics (MD) approach through force fields and GROMACS package is used to do MD simulations to locate sites with favorable interaction energy.

Keywords: GABA_A receptors, Lavender plant, Molecular dynamics, GROMACS

References

1. Kuffler SW, Edwards C (November 1958). "Mechanism of gamma aminobutyric acid (GABA) action and its relation to synaptic inhibition". *J. Neurophysiol.* **21** (6): 589–610.
2. Kravitz EA, Kuffler SW, Potter DD (September 1963). "Gamma-aminobutyric acid and other blocking compounds in crustacea. III. Their relative concentrations in separated motor and inhibitory axons". *J. Neurophysiol.* **26**: 739–51.
3. Krnjević K, Schwartz S (1967). "The action of gamma-aminobutyric acid on cortical neurones". *Exp Brain Res* **3** (4): 320–36.
4. Takeuchi A, Takeuchi N (August 1967). "Anion permeability of the inhibitory post-synaptic membrane of the crayfish neuromuscular junction". *J. Physiol. (Lond.)* **191** (3): 575–90.

5. Takeuchi A, Takeuchi N (November 1969). "A study of the action of picrotoxin on the inhibitory neuromuscular junction of the crayfish". *J. Physiol. (Lond.)* **205** (2): 377–91.

6. Mrs. M. Grieve, *A Modern Herbal*, Vol. II (New York: Dover Publications, Inc., 1971; ISBN 0-486-22799-5)

7. Kasper, S; Gastpar, M; Müller, WE; Volz, HP; Möller, HJ; Dienel, A; Schläfke, S (2010). "Silexan, an orally administered Lavandula oil preparation, is effective in the treatment of 'subsyndromal' anxiety disorder: a randomized, double-blind, placebo controlled trial". *International clinical psychopharmacology* 25 (5): 277–87. doi:10.1097/YIC.0b013e32833b3242. PMID 20512042

8. Inouye, S.; Takizawa, T.; Yamaguchi, H. (2001). "Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact". *Journal of Antimicrobial Chemotherapy* 47 (5): 565–73.